

European Journal of Vascular & Endovascular Surgery

Volume 48 • Issue 1 • 2014
CONTENTS

EDITORIAL

- 1 The Chronic Cerebrospinal Venous Insufficiency Debate
S. Onida and A.H. Davies

OBITUARY

- 4 Édouard Kieffer: A Master in Vascular Surgery
A. Branchereau and J.-B. Ricco

CAROTID DISEASE

- 5 Carotid Artery Atherosclerosis Among 65-year-old Swedish Men – A Population-based Screening Study
D. Högborg, B. Kragsterman, M. Björck, J. Tjärnström and A. Wanhainen
- 2.3% of 65-year-old males invited for AAA screening had a 50–99% stenosis or occlusion of a carotid artery, but only 40% were on antiplatelet or statin therapy.*

Invited Commentary

- 11 Commentary on ‘Carotid Artery Atherosclerosis Among 65-year-old Swedish Men: A Population Based Screening Study’
E.S. Debus

AORTIC DISEASE

- 13 **Editor's Choice** – Treatment Decisions for Descending Thoracic Aneurysm: Preferences for Thoracic Endovascular Aneurysm Repair or Surveillance in a Discrete Choice Experiment
CME *N. Rudarakanchana, B.C. Reeves, C.D. Bicknell, F.M. Heatley, N.J. Cheshire and J.T. Powell*

Amongst specialists, TEVAR was less likely to be preferred in patients aged >75 years, women, ASA Grade 4 and in patients requiring >25cm aortic coverage.

- 23 Monte Carlo Simulations for Assessment of Organ Radiation Doses and Cancer Risk in Patients Undergoing Abdominal Stent-graft Implantation
M.A. Blaszkak and R. Juszkat
- Radiation associated cancer risks with EVAR were low but slightly increased in younger patients and those where the neck angulation was >45 degrees.*

- 29 Parallel Endografts in the Treatment of Distal Aortic and Common Iliac Aneurysms
S. Lepidi, M. Piazza, P. Scrivere, M. Menegolo, M. Antonello, F. Grego and P. Frigatti
- Parallel endografts were an option in patients with distal aortic and bilateral CIA aneurysms, preserving flow into at least one internal iliac artery.*

- 38 Comparison of Three Contemporary Risk Scores for Mortality Following Elective Abdominal Aortic Aneurysm Repair
S.W. Grant, G.L. Hickey, E.D. Carlson and C.N. McCollum
- The British Aneurysm Repair Score delivered good procedural discrimination for predicting mortality after elective open and EVAR repair but needs to be regularly revalidated.*

Invited Commentary

- 45 Commentary on ‘Comparison of Three Contemporary Risk Scores for Mortality Following Elective Abdominal Aortic Aneurysm (AAA) Repair’
I. Loftus

CONTENTS—continued

- 46 Transbrachial and Femoral Artery Approach Endovascular Therapy for Flush Infrarenal Aortic Occlusion

L. Yuan, J. Bao, Z. Zhao, X. Feng, Q. Lu and Z. Jing

Experience of catheter-directed thrombolysis followed by transbrachial or femoral access endovascular treatment for flush infrarenal aortic occlusions.

- 53 Spontaneous Delayed Sealing in Selected Patients with a Primary Type-Ia Endoleak After Endovascular Aneurysm Repair

F. Bastos Gonçalves, H.J.M. Verhagen, K. Vasanthananthan, H.J.A. Zandvoort, F.L. Moll and J.A. van Herwaarden

All but one of 15 primary type-1a endoleaks detected intraoperatively and left untreated, sealed spontaneously, but one patient died following early rupture.

PERIPHERAL ARTERIAL DISEASE

- 60 Healthcare Quality Indicators of Peripheral Artery Disease Based on Systematic Reviews

S. Bellmunt, M. Roqué, D. Osorio, H. Pardo, J.-R. Escudero and X. Bonfill

Six quality indicators were defined based upon strong recommendations and methodological evidence for assessing appropriateness of delivering healthcare in PAD patients.

Invited Commentary

- 70 Surgeon Still a Step Behind Medicine: Indicators of Healthcare for Peripheral Artery Disease

P. De Rango

- 72 Healthcare-associated Infections After Lower Extremity Revascularization

A. Daryapeyma, O. Östlund and C.-M. Wahlgren

Swedvasc observed a 7% prevalence of soft tissue/wound infection after lower limb revascularization.

Invited Commentary

- 78 Commentary on “Healthcare-associated Infections after Lower Extremity Revascularization”

N. Chakfe, Y. Georg and D. Christmann

- 80 Radial Artery Bypass Graft Is a Feasible and Durable Conduit for Challenging Infrainguinal Revascularization: 17 Years of Melbourne Experience

B.M.E. Mees, D.R. Robinson, G. Fell and A.T.W. Chan

With an average length of 20cm, radial artery grafts provided an alternative autologous conduit source with a 5-year primary patency of 50%.

- 88 Systematic Review and Meta-analysis of Direct Versus Indirect Angiosomal Revascularisation of Infrapopliteal Arteries

D.C. Bosanquet, J.C.D. Glasbey, I.M. Williams and C.P. Twine

Direct angiosomal revascularization delivered improved wound healing and limb salvage compared to indirect revascularization.

VENOUS DISEASE

- 98 Role of Sinus in Prosthetic Venous Valve

W.-H. Tien, H.Y. Chen, Z.C. Berwick, J. Krieger, S. Chambers, D. Dabiri and G.S. Kassab

If a bicuspid, prosthetic venous valve is to be implanted without a sinus, a smaller valve leaflet size was preferable.

CORRESPONDENCE

- 105 Ocular Defects as Surrogate End-Points in Trials Comparing Carotid Endarterectomy and Stenting

A.D. Giannoukas, K.I. Paraskevas, R. Bulbulia and A. Halliday

- 106 Progressive compression, plantar load and calf venous pump

A. Caggiati and A. Mander

- 107 Response to ‘Re. Improvement of Venous Pumping Function by Double Progressive Compression Stockings: Higher Pressure Over the Calf is More Important Than a Graduated Pressure Profile’

G. Mosti

- 107 Re. 'Carotid Endarterectomy with Stent Removal in Management of In-stent Restenosis: A Safe, Feasible, and Effective Technique'
L. Davidović, I.B. Koncar, M. Markovic, N. Ilic, M. Dragas and D. Markovic
- 108 Response to 'Re. Carotid Endarterectomy with Stent Removal in Management of In-stent Restenosis: A Safe, Feasible, and Effective Technique'
J. Zheng and J. Zhao
- 109 Predictor Factors for Mortality in Hybrid Aortic Procedures
O. Gokalp, B. Eygi, Y. Besir, L. Yilik, I. Yurekli and A. Gurbuz
- 109 Re: 'Predictor Factors for Mortality in Hybrid Aortic Procedures'
S. Ben Ahmed and E. Rosset

MISCELLANEOUS

- 110 Selected Abstracts from the July Issue of the Journal of Vascular Surgery
- 114 Forthcoming Events

Editor's Choice: This paper has been selected by the Editor to be made available online as Open Access.



To access continuing medical education questions on this paper, please go to www.vascular education.com and click on 'CME'